

POSTER PRESENTATION

Open Access

Modeling spontaneous activity across an excitable epithelium: support for a coordination scenario of early neural evolution

Oltman O de Wiljes^{1*}, Ronald AJ van Elburg², Michael Biehl³, Fred A Keijzer¹*From* 24th Annual Computational Neuroscience Meeting: CNS*2015
Prague, Czech Republic. 18-23 July 2015

The reason why nervous systems first arose is an open question. Internal coordination models hold that nervous systems evolved initially as a device to coordinate internal activity, enabling multicellular effectors. They stress the use of multicellular contractility as an effector for motility: some sort of coordinative structure would have been necessary to have multicellular effectors in the first place. A recent example of such a model, the skin brain thesis, suggests that excitable epithelia using chemical signaling are a potential candidate as a nervous system precursor.

We developed a computational model and a measure for whole body coordination to investigate the coordinative properties of such excitable epithelia. Using this measure we show that excitable epithelia can spontaneously exhibit body-scale patterns of activation (see Figure 1.). Relevant factors determining the extent of patterning are the noise level for exocytosis, relative body dimensions, and body size. In smaller bodies whole-body coordination emerges from cellular excitability and bidirectional excitatory transmission alone.

Our results show that basic internal coordination as proposed by the skin brain thesis could have arisen in this potential nervous system precursor, providing support that this configuration may have played a role as a proto-neural system and requires further investigation.

Authors' details¹Department of Theoretical Philosophy, Groningen University, Groningen, the Netherlands. ²Institute of Artificial Intelligence, Groningen University,Groningen, the Netherlands. ³Johann Bernoulli Institute for Mathematics and Computer Science, Groningen University, Groningen, the Netherlands.

Published: 18 December 2015

doi:10.1186/1471-2202-16-S1-P120

Cite this article as: de Wiljes et al.: Modeling spontaneous activity across an excitable epithelium: support for a coordination scenario of early neural evolution. *BMC Neuroscience* 2015 **16**(Suppl 1):P120.**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit* Correspondence: o.o.de.wiljes@rug.nl¹Department of Theoretical Philosophy, Groningen University, Groningen, the Netherlands

Full list of author information is available at the end of the article




Figure 1 Wave patterns on a long tube (length: 32 cells, circumference: 8 cells, noise rate: 0.1 Hz). (A) Network geometry: the scissors indicate the line at which the tube is cut for presentation in **B**. **(B)** Snapshots of network activity during 4 ms intervals in an illustrative phase of the dynamics. The wave fronts propagating transversely collide with each other, causing extinction due to the refractory period. Subsequently, the remaining wave-fronts propagating longitudinally dominate the dynamics and the North-South wave-front orientation dominates. **(C)** Temporal development of different wave-front orientations. Snapshot time markings are consistent with those in **B**.